

Solving the Richardson equations close to the critical points

F. Domínguez¹, C. Esebbag¹ and J. Dukelsky²

¹ *Departamento de Matemáticas, Universidad de Alcalá, 28871 Alcalá de Henares, Spain.*

² *Instituto de Estructura de la Materia. CSIC. Serrano 123. 28006 Madrid. Spain.*

We study the Richardson equations close to the critical values of the pairing strength g_c where the occurrence of divergencies preclude numerical solutions. We derive a set of equations for determining the critical g values and the non-collapsing pair energies. Studying the behavior of the solutions close to the critical points, we develop a procedure to solve numerically the Richardson equations for arbitrary coupling strength.

PACS numbers: 02.30.Ik, 71.10.Li, 74.20.Fg

I. INTRODUCTION

The exact solution of the BCS or pairing Hamiltonian was presented by Richardson in series of papers beginning in 1963 [1, 2], just a few years after the seminal paper of Bardeen, Cooper and Schrieffer [3]. Despite of the new avenues of research that it could have opened at this early stage in the development of the theory of superconductivity, the work passed almost unnoticed due, perhaps, to the technical difficulties in solving numerically the set of non linear equations for the spectral parameters. In recent year, the Richardson exact solution was rediscovered and applied successfully to ultrasmall metallic grains where it was shown to be essential for the description the soft crossover between superconductivity and the pairing fluctuation regime as a function of the grain size [4]. Since then the Richardson model has been extended to a wide class exactly solvable models called Richardson-Gaudin (RG) models with potential applications to several quantum many body fermion and boson systems [5, 6, 7, 8]. However, the numerical treatment of the exact solution for moderate to large size systems is still a cumbersome task in spite of the recent efforts to overcome this problem.

For the sake of clarity, let us begin by introducing the Richardson equations for an M fermion pair system:

$$1 - 4g \sum_{j=1}^L \frac{d_j}{2\eta_j - e_\alpha} + 4g \sum_{\substack{\beta=1 \\ (\beta \neq \alpha)}}^M \frac{1}{e_\alpha - e_\beta} = 0, \quad (1)$$

where g is the pairing strength, η_j are a set of L parameters usually related to the single particle energies, d_j are the effective degeneracies defined as $d_j = \nu_j/2 - \Omega_j/4$ with ν_j de number of unpaired fermions in level j (Seniority quantum number) and Ω_j the total degeneracy of the level j . The e_α 's are the M unknowns parameters called pair energies. Given the L parameters η_j , the L effective degeneracies d_j and a pairing strength g , the pair energies e_α are obtained by solving the set of M nonlinear equations (1). However, it is in general not easy to find a good initial guess that would lead directly to the appropriate solution. Instead, it is customary to begin with a given configuration in the weak coupling limit ($g \rightarrow 0$) where the equations (1) can only be satisfied for $e_\alpha \rightarrow 2\eta_j$. The exact solution is then evolved step by step for increasing values of g up to the desired value [9]. In each step one uses the solution obtained in the previous step as the input data for the numerical solver. Even if the procedure is successful, leading to the correct solution, it involves a heavy numerical work. In most cases, this procedures is stopped due to the existence of singularities for some critical values g_c of the pairing strength. At $g = g_c$ a subset of pair energies e_α turn out to be equal to $2\eta_k$ for some k , giving rise to divergencies in some terms of eq. (1) [10]. Remarkably, these divergencies cancel out and the corresponding solution (the set of e_α) is a continuous function of g in a neighborhood of g_c . In fact, the solutions are always continuous for every value of g .

The existence of such critical points has significant consequences in the numerical solution of the equations which becomes unstable near g_c . Very slow convergency, no convergency at all or jumping to another solution are the typical problems found close to g_c . Moreover, there may be more than just one critical value in the real interval $[0, g]$, and we do not know *a priori* where the g_c 's are located. Those values are known only after one has already solved the equations for a large number of points surrounding the g_c 's and making some kind of interpolation afterward. Therefore, the localization of the critical values of g relies on heuristic procedures and is not based on any mathematical properties of the equations.

An alternative procedure to cross the critical region, based on a non linear transformation of the collapsing pair energies was recently proposed [11]. However, this procedure is unable to predict the critical values of g .

In this paper we will analyze the properties of Richardson equations in the vicinity of critical g values. We will derive conditions that allow to *a priori* determine all the critical values g_c associated to any "single particle level" η_j ,

and provide the exact solution at these points. In addition, we will describe the asymptotic behavior of the solution in the limit $g \rightarrow g_c$, and we will present an algorithm to solve the Richardson equations for values of g near any g_c .

II. TRANSFORMING THE RICHARDSON EQUATIONS. THE CLUSTER EQUATIONS

Our approach to deal with the singularities in eq. (1) consists in transforming the system through a change of variables first suggested by Richardson [10] and later used by Rombouts *et al.* [11] to develop a numerical algorithm for solving the equations near the critical points. We have already mentioned that for critical values of g some subset of the pair energies becomes equal to one of the values of $2\eta_k$. It can be shown [10] that the number of such pair energies is $M_k = 1 - 2d_k$. We can then characterize a critical point g_c by the condition $\lim_{g \rightarrow g_c} e_\alpha = 2\eta_k \forall \alpha \in C_k$, where C_k stands for the subset of indices of the M_k pair energies that satisfy that limit, i.e. the e_α 's that cluster around the real point $2\eta_k$ in the complex plane for g near g_c (see [12] for a graphical representation of these clusters). Therefore, we will deal with two subsets of variables, the M_k e_α 's with $\alpha \in C_k$ that give rise to the singularities, and the remaining $(M - M_k)$ variables with $\alpha \notin C_k$ which behave smoothly close to g_c . Consequently, we will treat separately the M_k Richardson equations with $\alpha \in C_k$ (the cluster equations):

$$1 - 4g \frac{d_k}{2\eta_k - e_\alpha} - 4g \sum_{\substack{j=1 \\ (j \neq k)}}^L \frac{d_j}{2\eta_j - e_\alpha} + 4g \sum_{\substack{\beta \in C_k \\ (\beta \neq \alpha)}} \frac{1}{e_\alpha - e_\beta} + 4g \sum_{\beta \notin C_k} \frac{1}{e_\alpha - e_\beta} = 0. \quad (2)$$

The second and fourth terms of eq. (2) diverge for $g \rightarrow g_c$ since $(2\eta_k - e_\alpha)$ and $(e_\alpha - e_\beta)$ go to zero. Moreover these quantities must approach zero at the same rate in order to cancel out. To avoid the singularities we will multiply the cluster equations by $(2\eta_k - e_\alpha)^p$ (for some $p > 0$). With this in mind we will introduce a change of variables for the pair energies in the cluster (see ref. [10, 11])

$$S_p = \sum_{\alpha \in C_k} (2\eta_k - e_\alpha)^p \quad \text{for} \quad p = 1, 2, \dots, M_k. \quad (3)$$

This is an invertible transformation and, in principle, we can always recover the e_α 's for any arbitrary set of S_p . Keeping in mind that we have just M_k independent variables S_p we will extend the definition allowing p to be any positive integer or zero (note that $S_0 = M_k$). The variables S_p behave smoothly in the vicinity of g_c , they are real and, for $p > 0$, they satisfy $\lim_{g \rightarrow g_c} S_p = 0$.

In order to transform the cluster equations (2) we first multiply them by $(2\eta_k - e_\alpha)^p$, for a generic integer power p , and then sum the resulting equations over $\alpha \in C_k$:

$$S_p + 2gp S_{p-1} - 2g \sum_{i=0}^{p-1} S_{p-i-1} S_i - 4g \sum_{\substack{j=1 \\ (j \neq k)}}^L \sum_{\alpha \in C_k} \frac{d_j (2\eta_k - e_\alpha)^p}{2\eta_j - e_\alpha} + 4g \sum_{\beta \notin C_k} \sum_{\alpha \in C_k} \frac{(2\eta_k - e_\alpha)^p}{e_\alpha - e_\beta} = 0. \quad (4)$$

Note that the last two terms in the left hand side of the equation cannot be easily rewritten in terms of S_p if we restrict to values of $p \leq M_k$. To overcome this limitation, we will make a series expansion of those terms, allowing p to be any integer. We are interested in the limiting behavior of the solution close to g_c so that our reasoning will be valid in an interval $g \in (g_c - \delta, g_c + \delta)$ for some small enough radius $\delta > 0$. Taking into account that $(2\eta_k - e_\alpha)$ is infinitesimal at g_c and that e_β in equation (4) does not belong to the cluster, we have:

$$\begin{aligned} \frac{1}{2\eta_j - e_\alpha} &= \frac{1}{2\eta_j - 2\eta_k} \sum_{n=0}^{\infty} \left(\frac{2\eta_k - e_\alpha}{2\eta_k - 2\eta_j} \right)^n, \\ \frac{1}{e_\alpha - e_\beta} &= \frac{1}{2\eta_k - e_\beta} \sum_{n=0}^{\infty} \left(\frac{2\eta_k - e_\alpha}{2\eta_k - e_\beta} \right)^n. \end{aligned} \quad (5)$$

These are absolutely (and rapidly) convergent geometric series, a property that will remain valid after a finite summation over α . Introducing eq. (5) in eq. (4) and making the summation over α we get for $p = 1$:

$$S_1 - 4g d_k M_k - 2g(M_k^2 - M_k) + 4g \sum_{n=0}^{\infty} S_{1+n} P_n = 0, \quad (6)$$

where the $\{P_n\}$ is a set of coefficients that will be defined below. Taking in (6) the limit $g \rightarrow g_c$ the second and third terms must cancel out and one gets the cluster condition $M_k = 1 - 2d_k$. Similarly, for $p > 1$ we obtain:

$$S_p - 2g(M_k + 1 - p)S_{p-1} - 2g \sum_{i=1}^{p-2} S_{p-i-1}S_i + 4g \sum_{n=0}^{\infty} S_{p+n}P_n = 0 \quad \text{for all } p > 1. \quad (7)$$

The coefficients P_n appearing in (6) and (7) are defined as:

$$P_n = \sum_{\substack{j=1 \\ (j \neq k)}}^L \frac{d_j}{(2\eta_k - 2\eta_j)^{n+1}} + \sum_{\beta \notin C_k} \frac{1}{(2\eta_k - e_\beta)^{n+1}}. \quad (8)$$

The infinite recurrence relation (7) is the fundamental equation of our formalism. In order to achieve our main goals we need to establish some important properties (lemma 1 and lemma 2) of the variables S_p and the series appearing in eqs. (6) and (7). It is easy to show that these series are bounded by their first element S_p through the condition:

$$\left| \sum_{n=0}^{\infty} S_{p+n}P_n \right| \leq |S_p| D \frac{1}{1-x}, \quad (9)$$

where D is a positive bounded number involving η_j and e_β , and x is an upper bound for the absolute values of the geometric series ratios in eq.(5) satisfying $\lim_{g \rightarrow g_c} x = 0$. On the other hand, we have that $|S_p| \leq \sum_{\alpha \in C_k} |2\eta_k - e_\alpha|^p$ with $(2\eta_k - e_\alpha) \rightarrow 0$ as $g \rightarrow g_c$. Therefore, we will assume in what follows that there is an integer number n_δ big enough such that we may disregard every term S_p (and related series) for $p > n_\delta$.

The next aspect we need to analyze is the order of the infinitesimal S_p in relation to the order of S_1 . We will demonstrate that $S_1, S_2, S_3, \dots, S_{M_k}$ are of the same order, implying that $\lim_{g \rightarrow g_c} S_p/S_1 = \chi_p \neq 0$ for $p \leq M_k$. On the other hand, S_p is of higher order than S_1 if $p > M_k$, that is $\lim_{g \rightarrow g_c} S_p/S_1 = 0$. In lemmas 1 and 2 we will demonstrate both assertions.

Lemma 1. *Assuming that $(1 + 4gP_0) \neq 0$, with $g \in (g_c - \delta, g_c + \delta)$ and P_0 defined in (8), the set $\{S_2, S_3, \dots, S_\mu\}$, for some positive integer μ with $2 \leq \mu \leq M_k$, are infinitesimal of the same order as S_1 at $g = g_c$. In the general case $\mu = M_k$.*

Proof. Rearranging equation (7) we get:

$$S_p = \frac{1}{(1 + 4gP_0)} \left[2g(M_k + 1 - p)S_{p-1} + 2g \sum_{i=1}^{p-2} S_{p-i-1}S_i - 4g \sum_{n=1}^{\infty} S_{p+n}P_n \right]. \quad (10)$$

To prove our statement we will apply this equation recursively starting with $p = 2$. In this case we get:

$$S_2 = \frac{2g(M_k - 1)}{(1 + 4gP_0)} S_1 - \frac{4g}{(1 + 4gP_0)} \sum_{n=1}^{\infty} S_{2+n}P_n,$$

so that S_2 is of the same order as S_1 . In the next step we set $p = 3$ in eq. (10), getting S_3 in terms of S_2 and S_p 's with $p > 3$. Replacing S_2 with the value obtained previously and rearranging the resulting expression we get for S_3 :

$$S_3 = \frac{1}{(1 + 4gP_0)^2 + 8g^2P_1(M_k - 2)} \left[(2g)^2(M_k - 2)(M_k - 1)S_1 + 2g(1 + 4gP_0)S_1^2 - 4g \sum_{n=1}^{\infty} (2g(M_k - 2)P_{n+1} + (1 + 4gP_0)P_n)S_{3+n} \right],$$

from where we see that S_3 is of the same order as S_1 . We can continue this process replacing in each step S_{p-1} in eq. (10) for the expression computed in the previous step and solving again for S_p , getting for every $p > 2$ and $p \leq M_k$:

$$S_p = B_p S_1 + \sum_{n=1}^{\infty} D_n^{(p)} S_{p+n} + \sum_{\substack{i,j \\ i+j < p}} X_{i,j}^{(p)} S_i S_j \quad (11)$$

where the coefficients B_p , $D_n^{(p)}$ and $X_{i,j}^{(p)}$ can be calculated recursively. For instance, for the linear terms we have the recurrence relation

$$\begin{aligned} B_{p+1} &= \frac{2g(M_k - p)}{(1 + 4gP_0) - 2g(M_k - p)D_1^{(p)}} B_p, \\ D_n^{(p+1)} &= \frac{2g}{(1 + 4gP_0) - 2g(M_k - p)D_1^{(p)}} \left((M_k - p)D_{n+1}^{(p)} - 2P_n \right), \end{aligned} \quad (12)$$

starting with the known coefficients for $p = 2$.

In this way we have shown that the local expansion of S_p has linear terms in S_1 and in S_{p+n} plus quadratic terms, thus we conclude that S_p is an infinitesimal of the same order as S_1 for $1 \leq p \leq M_k$.

Let us now consider what happens when p takes the value $M_k + 1$. In such a case the factor $(M_k + 1 - p)$ in eq. (10) becomes zero and the chain of replacement is broken. The expansion of S_{M_k+1} has only linear terms of higher order and quadratic terms involving S_1 . Therefore for $p > M_k$ there are no linear terms in S_1 . We could have reached the same conclusion analyzing the recurrence relation (12).

Finally and for the sake of completeness, let us mention that after replacing S_{p-1} in equation (10) the terms involving S_p on both sides of the equation might cancel out. It would be equivalent to the cancellation of the denominators in the recurrence relation (12). In such a case our process of replacement is interrupted and ends for $p = \mu < M_k$. $\square$

So far, we have shown that the first S_p 's are of the same order as S_1 . At the same time, it was suggested that S_p is of higher order if $p > M_k$. In the next lemma we will proof this statement.

Lemma 2. *The set of variables $\{S_p\}$ for $p > M_k$ are infinitesimal of higher order than S_1 at $g = g_c$.*

Proof. In this case we proceed in the reverse way as we have done in lemma 1. We will begin with an S_p for a large value of p and carry out the replacement in backward direction. Once again we start rewriting equation (7) to obtain:

$$S_p = \frac{1}{(M_k - p)} \left[\frac{(1 + 4gP_0)}{2g} S_{p+1} + 2 \sum_{n=1}^{\infty} S_{p+1+n} P_n - \sum_{i=1}^{p-1} S_{p-i} S_i \right]. \quad (13)$$

We will now take, for $g \in (g_c - \delta, g_c + \delta)$, an integer number n_δ large enough such that we may disregard every term S_p for $p > n_\delta$. We start making $p = n_\delta$ in eq. (13). After removing the negligible terms it leads us to:

$$S_{n_\delta} = \frac{1}{(M_k - n_\delta)} \sum_{i=1}^{n_\delta-1} S_{n_\delta-i} S_i.$$

In the second step we make $p = n_\delta - 1$ in eq. (13) and replace the value of S_{n_δ} computed before. Repeating the same operation for $p = n_\delta - 2, n_\delta - 3, \dots, M_k + 1$ and replacing, in each case, the linear term for its value computed in the previous step, eventually we get the following quadratic expression for every $p > M_k$:

$$S_p = \sum_{\substack{i,j \\ i+j \geq p}} Y_{i,j}^{(p)} S_i S_j. \quad (14)$$

In this equation the coefficients $Y_{i,j}^{(p)}$ can be calculated recursively and the indices i, j take values starting from 1 but keeping the condition $i + j \geq p$. As only quadratic terms are retained we conclude that S_p for $p > M_k$ are infinitesimal of higher order than $S_1, S_2, \dots, S_{M_k}$. Note once again that the chain of replacement cannot be continued for $p \leq M_k$. $\square$

So far we have shown some essential properties of the variables S_p and the related series that are required to develop our formalism. In the next section we will derive the equations that allow to calculate the values of g_c associated with any single particle level η_k .

III. CRITICAL VALUES OF g AND THE SOLUTION OF RICHARDSON EQUATIONS

In this section we will derive a set of equations suitable to compute all the values of g_c for any single particle level η_k . Furthermore, we will obtain, at the same time, the solution of Richardson equation at g_c , namely, the values of the unknowns e_β for $\beta \notin C_k$, assuming that the variables in the cluster reach their limiting value $e_\alpha = 2\eta_k$ ($e_\alpha \in C_k$). We summarize our results in the following theorem:

An appropriate set of starting values at $g_0 = g_c + \delta g$ can be obtained by means of a linear approximation for the parameters S_p and e_α :

$$\begin{aligned} S_1(g_0) &\approx \left(\frac{dS_1}{dg} \right)_{g=g_c} \delta g, & S_p(g_0) &\approx \left(\frac{dS_1}{dg} \right)_{g=g_c} \chi_p \delta g \quad (p > 1), \\ e_\alpha(g_0) &\approx e_\alpha(g_c) + \left(\frac{de_\alpha}{dg} \right)_{g=g_c} \delta g \quad (\alpha \notin C_k), \end{aligned} \quad (18)$$

this approximate values will be the initial guess to solve the Richardson equations.

In order to determine the derivative dS_1/dg at $g = g_c$ we will consider up to second order terms in equations (6) and (7) and substitute the variables S_p in terms of the quadratic expression $S_p \approx \chi_p S_1 + a_p S_1^2$, where $\chi_p = 0$ for $p > M_k$, $\chi_1 = 1$ and $a_1 = 0$. From equation (14) we can see that S_p has quadratic terms in S_1 only for $p \leq 2M_k$, while for greater values of p the order in S_1 is higher due to the condition $i + j \geq p$. Therefore, we will retain just the first $2M_k$ variables S_p and deal with a system of $2M_k$ equations. Computing the derivative with respect to g of the resulting system, and after making some simple algebraic manipulations, we get for $p = 1$:

$$4g \sum_{n=1}^{2M_k} [(\chi_n + a_n S_1) P'_{n-1} + (a_n P_{n-1}) S'_1] = \frac{1}{g}, \quad (19)$$

and for $1 < p \leq 2M_k$:

$$\begin{aligned} &S'_1 \left[a_p - 2g(M_k + 1 - p)a_{p-1} - 2g \sum_{i=1}^{p-2} (\chi_{p-i-1} \chi_i) \right] + \\ &4g \sum_{n=0}^{2M_k-p} [(\chi_{n+p} + a_{n+p} S_1) P'_n + a_{n+p} P_n S'_1] = \frac{\chi_p + a_p S_1}{g}, \end{aligned} \quad (20)$$

where the primes stand for the derivatives with respect to g and, according to definition (8), we have:

$$P'_n = \sum_{\beta \notin C_k} \frac{(n+1)}{(2\eta_k - e_\beta)^{n+2}} \left(\frac{de_\beta}{dg} \right). \quad (21)$$

Taking the limit $g \rightarrow g_c$ and reordering the equations in a convenient way we obtain

$$\left(-\frac{1}{g_c} + 4g_c \sum_{n=1}^{M_k} \chi_n P'_{n-1} \right) + \sum_{n=2}^{2M_k} (4P_{n-1})(a_n S'_1) = 0, \quad (22)$$

for $p = 1$, and

$$\begin{aligned} &\left(-\frac{\chi_p}{g_c} - 2g_c S'_1 \sum_{i=1}^{p-2} (\chi_{p-i-1} \chi_i) + 4g_c \sum_{n=0}^{M_k-p} \chi_{n+p} P'_n \right) - 2g_c(M_k + 1 - p) a_{p-1} S'_1 + \\ &(1 + 4g_c P_0) a_p S'_1 + \sum_{j=p+1}^{2M_k} (4g_c P_{j-p})(a_j S'_1) = 0, \end{aligned} \quad (23)$$

for $1 < p \leq 2M_k$. This is a non-homogeneous system of $2M_k$ equations with a set of $2M_k + (M - M_k)$ unknowns: S'_1 , $a_2, a_3, \dots, a_{2M_k}$ joined with the derivatives (de_β/dg) , for $\beta \notin C_k$.

The system of equations (23) is non-linear because of the products $a_j S'_1$. However, we can obtain a linear system for the derivatives with a unique solution by defining a matrix $\mathbf{B}$ as:

$$\begin{aligned} B_{p,1} &= \left(-\frac{\chi_p}{g_c} - 2g_c S'_1 \sum_{i=1}^{p-2} (\chi_{p-i-1} \chi_i) + 4g_c \sum_{n=0}^{M_k-p} \chi_{n+p} P'_n \right), \\ B_{p,p-1} &= -2g_c(M_k + 1 - p), & B_{p,p} &= (1 + 4g_c P_0) \quad (p > 1), \\ B_{p,j} &= 4g_c P_{j-p} \quad (j > p), & B_{p,j} &= 0 \quad (1 < j < p-1); \end{aligned}$$

and the non-null vector $\mathbf{v}$:

$$\mathbf{v} = \left(1, a_2 \frac{dS_1}{dg}, a_3 \frac{dS_1}{dg}, \dots, a_{2M_k} \frac{dS_1}{dg} \right).$$

Using these definitions, equations (23) can be rewritten as:

$$\mathbf{B} \cdot \mathbf{v} = \mathbf{0},$$

such that the following condition must be satisfied:

$$\det(\mathbf{B}) = 0. \quad (24)$$

Since the derivatives appear just in the first column of the matrix $\mathbf{B}$, (24) is a linear equation in $\frac{dS_1}{dg}$ and in the $(M - M_k)$ derivatives $\frac{de_\beta}{dg}$ (for $\beta \notin C_k$) that does not include the unknown coefficients a_p . A set of $(M - M_k)$ equations is required to complete the system. In order to get the necessary equations we will compute the derivatives of the Richardson equations out of the cluster:

$$1 - 4g \sum_{j=1}^L \frac{d_j}{2\eta_j - e_\alpha} + 4g \sum_{\substack{\beta \notin C_k \\ (\beta \neq \alpha)}} \frac{1}{e_\alpha - e_\beta} + 4g \sum_{\beta \in C_k} \frac{1}{e_\alpha - e_\beta} = 0 \quad (\alpha \notin C_k). \quad (25)$$

Expanding the last term in the variables S_p we obtain

$$1 - 4g \sum_{j=1}^L \frac{d_j}{2\eta_j - e_\alpha} + 4g \sum_{\substack{\beta \notin C_k \\ (\beta \neq \alpha)}} \frac{1}{e_\alpha - e_\beta} - 4g \sum_{n=0}^{\infty} \frac{1}{(2\eta_k - e_\alpha)^{n+1}} S_n = 0. \quad (26)$$

Computing the derivative with respect to g and taking the limit $g \rightarrow g_c$ we finally get:

$$\begin{aligned} & -\frac{1}{g_c} - 4g_c \sum_{j=1}^L \frac{d_j}{(2\eta_j - e_\alpha)^2} \left(\frac{de_\alpha}{dg} \right)_{g=g_c} + 4g_c \sum_{\substack{\beta \notin C_k \\ (\beta \neq \alpha)}} \frac{1}{(e_\alpha - e_\beta)^2} \left(\frac{de_\beta}{dg} - \frac{de_\alpha}{dg} \right)_{g=g_c} \\ & - 4g_c \frac{M_k}{(2\eta_k - e_\alpha)^2} \left(\frac{de_\alpha}{dg} \right)_{g=g_c} - 4g_c \left(\sum_{n=1}^{M_k} \frac{\chi_n}{(2\eta_k - e_\alpha)^{n+1}} \right) \frac{dS_1}{dg} = 0 \quad (\alpha \notin C_k). \end{aligned} \quad (27)$$

Equations (24) and (27) constitute a non-homogenous linear system with a unique solution for the desire derivatives.

V. A NUMERICAL EXAMPLE: FERMIONS IN A 2-DIMENSIONAL LATTICE

In this section we will apply our formalism to a pairing model of fermions in a 2-D square lattice of $N \times N$ sites with periodic boundary conditions. This example was previously treated in Ref. [5] for the ground state with a repulsive pairing interaction ($g > 0$). We will now analyze the model using the methodology developed here, focussing on the critical g values and in the behavior of the solutions in their vicinities.

We will consider a 6×6 lattice with 36 fermions (half filling, $M = 18$). The pairing Hamiltonian in momentum space is

$$H_P = \sum_{\mathbf{k}} \varepsilon_{\mathbf{k}} a_{\mathbf{k}}^\dagger a_{\mathbf{k}} + \frac{g}{2} \sum_{\substack{\mathbf{k} \\ \bar{\mathbf{k}}'}} a_{\mathbf{k}}^\dagger a_{\bar{\mathbf{k}}}^\dagger a_{\bar{\mathbf{k}}'} a_{\mathbf{k}'}$$

where $\mathbf{k} \equiv [k_x, k_y]$ and $\varepsilon_{\mathbf{k}} = -2[\cos(k_x) + \cos(k_y)] = \eta_{\mathbf{k}}$, and $\bar{\mathbf{k}}$ is the time reversal of $\mathbf{k}$. The single fermion energies ε_j and the corresponding degeneracies Ω_j for the 6×6 lattice are displayed in the Table I. In the limit $g = 0$ the ground state is obtained by distributing the $M = 18$ pairs in the lowest possible states. In this numerical example we will work within the fully paired subspace containing the ground state ($\nu_j = 0$, $d_j = -\Omega_j/4$). Consequently, the excited states to which we will refer to in this example are within the seniority 0 subspace.




FIG. 1: Real part of the pair energies e_α for attractive ($g < 0$) and repulsive ($g > 0$) pairing in a 6×6 lattice.

TABLE I: Single fermion energies and degeneracies for the 6×6 lattice.

j	1	2	3	4	5	6	7	8	9
ε_j	-4	-3	-2	-1	0	1	2	3	4
Ω_j	2	8	8	8	20	8	8	8	2

We will begin solving the system of equations (15) for a specific single particle level, for instance, $j = 4$ and $\varepsilon_4 = -1$ (see table I). The degeneracy of this level is 8 and the number of pair energies in the cluster that collapses at $2\varepsilon_4 = -2$ is given by the cluster condition $M_4 = 1 - 2d_4 = 5$, one more than the value allowed by the Pauli exclusion principle. The solution of the system (15) provides all the information concerning the many body state at each specific critical value of g . In table II we show, for $j = 4$, the first negative g_c 's together with the corresponding energy eigenvalues.

A different kind of analysis is presented in Table III where we show the first critical g values for the ground state. In this case the clusters collapse at different single particle levels for each g_c as indicated. All the cluster are formed by 5 pair energies except for $j = 1$ where $M_1 = 3$. In order to have a global vision of the results we show in Fig. 1 the real part of the pair energies for the ground state solution of the Richardson equations (1) for positive and negative g values. The full lines are the pair energies e_α and the horizontal dotted lines correspond to $2\varepsilon_j$. The critical points of Table III can be seen in the figure at the crossing point of each cluster with twice the corresponding single particle energy.

The solutions near the critical points g_c were obtained following the approach described in Section IV. With the solution for the first g_c next to $g = 0$ (negative or positive depending on the case), i.e, the e_β for $\beta \notin C_k$ and value of g_c itself, we solve the linear system of equations (24) and (27) to obtain the derivatives (dS_1/dg) and (de_β/dg) at the critical point. In addition we solve the linear system (16) to get the coefficients χ_p . With all this information we determine a good initial guess at some g_0 near g_c by mean of the linear approximation (18). Next we solve the Richardson equations at g_0 , a process that rapidly converge to an accurate numerical solution. With the solution at g_0 as a starting point we move to the next value of $g = g_0 + \Delta g$. Proceeding in this way and updating the starting values at each step we compute all the desired points in the curves. We repeat the process for each critical g . The pair energies obtained from the solutions for $g \approx 0$ smoothly connect with those obtained near the first g_c . The same smooth behavior is observed between two consecutive critical values.

The good quality of the initial guess near g_c is due to the smooth behavior of the variables S_p and e_β ($\beta \notin C_k$) that allows us to use the linear approximation (18). For the e_β 's this fact can be appreciated in figure 1. In figure 2 we show a graphical representation of the first S_p 's in a range of g around $g_c = 0.170878$ corresponding to the collapse at the single particle level $j = 2$, $2\varepsilon_2 = -6$ of a cluster of $M_2 = 5$ pair energies. We can see in the figure the linearity of the S_p 's for $1 \leq p \leq 5$ confirming that S_p is an infinitesimal of the same order as S_1 for $2 \leq p \leq M_2$. In addition, we have plotted the results for S_6 , which clearly shows that it is an infinitesimal of a higher order.




FIG. 2: Variables S_p close to the collapse at $g_c = 0.170878$. The first 5 S_p 's are displayed in solid lines while S_6 is drawn in dashed line.

TABLE II: First critical g values for the cluster collapsing at $2\varepsilon_4 = -2$

g_c	Energy
-0.0384565	-47.6184
-0.0391412	-49.5405
-0.0394719	-53.3549
-0.0404240	-55.5262
-0.0412922	-57.4106
-0.0413245	-62.5795

VI. CONCLUSIONS

In this work we have studied the solution of the Richardson equations close to the critical values of the coupling constant g . We have derived a set of well behaved equations to determine the actual critical values g_c associated with any single particle energy ε_k , and the complete set of pair energies defining the exact eigenstate of the system. In addition, we studied the behavior of the solutions close to the critical points, obtaining a linear approximation for the pair energies that serves as a good initial guess in the critical region. With this formalism, one can solve numerically the Richardson equations around the critical points, overcoming the numerical instabilities that usually arise. The knowledge of the critical values of the coupling constant greatly simplifies the numerical treatment of the equations. To illustrate the formalism we have analyzed the pairing Hamiltonian in a 6×6 square lattice at half filling for repulsive and attractive pairing strength. Making use of our approach we have located the g_c 's for the ground state and several excited states in the Seniority 0 subspace. The numerical solution between consecutive values of g_c 's can be easily carried out by solving the original Richardson equations (1). With this new approach we hope to be able to solve exactly large systems with arbitrary single particle energies and degeneracies.

TABLE III: Critical g values for the ground state

	$2\varepsilon_4 = -2$	$2\varepsilon_3 = -4$	$2\varepsilon_2 = -6$	$2\varepsilon_1 = -8$
g_c (negative)	-0.0413245	-0.0635021	-0.0877434	-0.131927
g_c (positive)	0.598232	0.240579	0.170878	—

acknowledgments

We acknowledge fruitful discussions S. Rombouts. This work was supported by Spanish DGI under grant BFM2003-05316-C02-02.

-
- [1] Richardson R W 1963 *Phys. Lett.* **3** 277.
 - [2] Richardson R W and Sherman N 1964 *Nucl. Phys.* **52** 221.
 - [3] Bardeen J, Cooper L N and Schrieffer J R 1957 *Phys. Rev.* **108** 1175.
 - [4] Dukelsky J and Sierra G 1999 *Phys. Rev. Lett.* **83** 172; Sierra *et al.* 2000 *Phys. Rev. B* **61** R11890.
 - [5] Dukelsky J, Esebbag C and Schuck P 2001 *Phys. Rev. Lett.* **87** 066403.
 - [6] Dukelsky J, Pittel S and Sierra G 2004 *Rev. Mod. Phys.* **76** 643.
 - [7] Links J *et al.* 2003 *J. Phys. A: Math. Gen.* **36** R63.
 - [8] Ortiz G *et al.* 2005 *Nucl. Phys. B* **707** 421.
 - [9] Richardson R W 1966 *Phys. Rev.* **141** 949.
 - [10] Richardson R W 1965 *J. Math. Phys.* **6** 1034.
 - [11] Rombouts S, Van Neck D and Dukelsky J 2004 *Phys. Rev. C* **69** 061303.
 - [12] Dukelsky J, Esebbag C and Pittel S 2004 *Phys. Rev. Lett.* **88** 062501.